

This paper appeared in the European J. of Neurosci., 1999, 11, 1891-1898.

Social intelligence in the normal and autistic brain: an fMRI study

Simon Baron-Cohen*, Howard A. Ring+, Sally Wheelwright*, Edward T.
Bullmore++, Mick J. Brammer++, Andrew Simmons**, and Steve C.R. Williams**.

*Departments of Experimental Psychology and Psychiatry,
University of Cambridge, Downing St., Cambridge, CB2 3EB, UK.

+Academic Department of Psychological Medicine,
St. Bartholomew's and the Royal London School of Medicine,
Whitechapel, London, E1 1BB, UK.

** Neuroimaging Research, Department of Clinical Neuroscience, and

++ Department of Biostatistics and Computing, Institute of Psychiatry,
University of London, Denmark Hill, London, SE5 8AF, UK.

Acknowledgments: This work was funded by a grant to SBC, HR, and SCW from the Wellcome Trust, and by a grant to the first author from the Gatsby Foundation. EB is also supported by the Wellcome Trust. We are grateful to Barry Everitt for comments on the first draft of this paper, and Chris Andrew for technical support. Correspondence and requests for materials to SBC (email: sb205@cus.cam.ac.uk).

Abstract

There is increasing support for the existence of “social intelligence”(Humphrey, 1984), independent of general intelligence. Brothers (Brothers, 1990) proposed a network of neural regions that comprise the “social brain”: the orbito-frontal cortex (OFC), superior temporal gyrus (STG) and amygdala. We tested Brothers’ theory by examining both normal subjects as well as patients with high-functioning autism or Asperger Syndrome (AS), who are well known to have deficits in social intelligence, and perhaps deficits in amygdala function (Bauman & Kemper, 1988). We used a test of judging from the expressions of another person’s eyes what that other person might be thinking or feeling. Using functional magnetic resonance imaging (fMRI) we confirmed Brothers’ prediction that the STG and amygdala show increased activation when using social intelligence. Some areas of prefrontal cortex also showed activation. In contrast, patients with autism or AS activated the fronto-temporal regions but not the amygdala when making mentalistic inferences from the eyes. These results provide support for the social brain theory of normal function, and the amygdala theory of autism.

Social intelligence encompasses our abilities to interpret others' behaviour in terms of mental states (thoughts, intentions, desires, and beliefs), to interact both in complex social groups and in close relationships, to empathize with others' states of mind, and to predict how others will feel, think, and behave. The idea that social intelligence might be independent of, **or dissociable from**, general intelligence comes from several sources: First, individuals exist who are capable of considerable understanding of the non-social world (e.g., physics, maths, engineering) yet who readily admit to finding the social world confusing (Baron-Cohen, Wheelwright, Stone & Rutherford, submitted; Sacks, 1994). The opposite type of individual also exists: people who have no difficulty interacting with the social world but who find non-social problem-solving confusing (Karmiloff-Smith, Grant, Bellugi & Baron-Cohen, 1995). Secondly, certain kinds of brain damage can cause selective impairment in social judgment (Damasio, Tranel & Damasio, 1990) without any necessary loss to general problem-solving ability. Loss of social judgment can co-occur with memory and executive dysfunction, following amygdala damage (Tranel & Hyman, 1990), but the functional double dissociation between social and non-social intelligence implies their neural independence. Finally, many primatologists now believe that social problem-solving was a key driving force behind the evolution of primate intelligence, rather than tool-use or other non-social problem solving (Whiten, 1991).

A neural basis of social intelligence was first proposed by Brothers (Brothers, 1990). She suggested from both animal lesion studies (Kling & Brothers, 1992), single cell recording studies (Brothers, Ring & Kling, 1990), and neurological studies (cited above) that this involves the amygdala, OFC, and STG. Together, she postulated these

comprise the “social brain”. Damage to the amygdala impairs judgment of emotion (Calder et al., 1996), damage to the OFC impairs judgement of what is socially appropriate (Eslinger & Damasio, 1985), and damage to the STG impairs face-perception (Campbell, Heywood, Cowey, Regard & Landis, 1990). Single cell recording studies in non-human primates also confirm the role of the STG in detection of gaze (Perrett et al., 1985). Recent PET and SPECT studies of “theory of mind” (or the ability to impute mental states) also implicate areas of prefrontal cortex, specifically the medial frontal cortex (MFC) (Fletcher et al., 1995; Goel, Grafman, Sadato & Hallett, 1995) and the OFC (Baron-Cohen et al., 1994).

The present fMRI study had 2 main aims: (1) To test Brothers’ social brain theory that these neural regions, identified independently from several different studies and methods of investigation, are jointly activated in a group of normal subjects performing a novel social intelligence test. (2) To test the validity of this neural model of social intelligence by comparing normal cerebral blood oxygenation changes induced by performance of this task to hypothetically abnormal changes in a group of patients with high-functioning autism or Asperger Syndrome, known to have social impairment (Baron-Cohen & Ring, 1994). In particular, we predicted abnormal amygdala activation in the autism¹ group, on the basis of 5 lines of evidence: (a) a neuroanatomical study of autism at post-mortem found microscopic pathology (in the form of increased cell density) in the amygdala, in the presence of normal amygdala volume (Bauman & Kemper, 1994; Rapin & Katzman, 1998); (b) the only animal model of autism involves ablation of the amygdala (in rhesus monkeys) (Bachevalier,

¹ For brevity here, we refer to the autism group, but remind the reader that this includes patients with high functioning autism.

1991). **Whilst there is some dispute as to whether one can have an animal model of autism when the syndrome involves deficits in higher order cognition, this is at least consistent with the amygdala theory;** (c) patients with amygdala lesions show impairments in social judgement (Adolphs, Tranel, Damasio & Damasio, 1994; Young, Hellawell, De Wal & Johnson, 1996)[(d) using SPECT, patients with autism spectrum conditions show significant reductions in temporal lobe blood flow, regardless of whether they have temporal lobe epilepsy (Gillberg, Bjure, Uvebrant, Vestergren & Gillberg, 1993); and (e) in cases of tuberous sclerosis, autistic comorbidity is determined by hamartomata in the temporal lobe (Bolton & Griffiths, 1997). For all these reasons, a basic impairment of amygdala function in autism seems very plausible².

Subjects

6 subjects with autism (4 m, 2 f) were matched for mean age, handedness, IQ, socioeconomic status, and educational level, with 12 subjects in the normal group (6m, 6f). IQ was assessed with the full Wechsler Adult Intelligence Scale (WAIS-R). Subjects were only included if their IQ was in the normal range (i.e., above 85 both in terms of full-scale IQ, and in terms of performance and verbal IQ). These variables are shown in Table 1. There were no significant differences on any of these dimensions. Individuals in the clinical group all had a diagnosis of autism or Asperger Syndrome, using DSM-IV (APA, 1994) and ICD-10 (1994) criteria.

² We emphasize the amygdala theory of autism, and it might be thought that this is too narrow, since some of the lines of evidence cited here implicate temporal lobe structures more widely, which include the amygdala but also include other adjacent mesiotemporal areas. To the extent that the results reported later support the amygdala theory, it remains for future work to establish the specificity of this finding.

insert Table 1 here

Experimental Design

We used a blocked periodic ABA... design. Each epoch (A or B) was presented for 30 secs, and there were 5 cycles of AB alternation in total. Images were acquired from each subject during visual presentation of two tasks, both of which involved deriving socially relevant information from facial stimuli. This periodically designed (ABA..) experiment was expected to induce periodic MR signal change with signal maximum during task A in brain regions relatively specialized for gender recognition from facial stimuli; and periodic MR signal change with signal maximum during task B in brain regions relatively specialized for mental state recognition from facial stimuli. The response involved a forced choice between the two words offered (pressing one of two buttons with the right hand to select the right or left word). Correct words were counterbalanced to left and right side. **Since both tasks were social, either may have resulted in anomalous activation in the autism group, though we predicted abnormalities would only arise in Task B.**

Method

Task A: Subjects were visually presented with a series of photographs of eyes and asked to indicate by right handed button press whether each stimulus was a man or a woman. In this first task (A: gender recognition), instructions to subjects were to

decide for each stimulus which of two simultaneously presented words (“male” or “female”) best described the face. Each stimulus was presented for 5 secs and was followed by a 0.75 sec interval in which the screen was blank. Stimuli were drawn from 30 faces of women or men. Stimuli were presented 3.5m from the subject, subtending visual angles of 10 degrees horizontally and 8 degrees vertically.

Task B: Subjects were presented with exactly the same stimuli but were asked to indicate by button press which of two simultaneously presented words best described the mental state of the photographed person. Thus, the key difference between the two tasks was the type of judgement the subject had to make when viewing the eyes³. Subjects were presented with an example of the stimuli before scanning. For this second task (B: theory of mind), instructions to subjects was to decide for each stimulus which of two simultaneously presented words best described what the person in the photograph was feeling or thinking. Task B is an “advanced” theory of mind test, in that it is used with adults.

Adults with high-functioning autism or AS, with intelligence in the normal range, show deficits on this task (Baron-Cohen, Jolliffe, Mortimore & Robertson, 1997), as do parents of children with autism/AS (Baron-Cohen & Hammer, 1997). Children with William’s Syndrome are not impaired on this test, despite their general retardation (Tager-Flusberg, Boshart & Baron-Cohen, 1998). Examples of the eyes used in the experimental condition, together with the forced choice words that

³ A secondary difference between tasks A and B is that in A the same words (male, female) always appear, whilst in B different words (describing a range of mental states) appear. This is inevitable if one uses the same pictorial stimuli in both tasks, whilst varying the social judgement required. However, we cannot see any reason why this factor should explain the results.

appeared underneath each face, are shown in Figure 1. **Finally, as a control pre-test outside the scanner, subjects were given the opportunity to pick out any words in a list of mental state words that would appear in Task B that they did not recognize or understand, in which case a glossary definition was provided by the experimenter. Neither group made use of this, reflecting that the words used were relatively common, and that the adult subjects in both groups were of normal intelligence.**

insert Figure 1 here

Image Acquisition and Analysis

Single shot gradient echo, echoplanar images were acquired using a 1.5 Tesla GE Signa system (General Electric, Milwaukee, WI, USA) fitted with Advanced NMR hardware and software (ANMR, Woburn, MA, USA) using a standard head coil. 100 T_2^* -weighted images depicting BOLD contrast (Ogawa, Lee, Kay & Tank, 1990) were acquired over 5 min at each of 14 near-axial non-contiguous 7-mm-thick planes parallel to the intercommissural (AC-PC) line, providing whole-brain coverage: TE, 40ms; TR, 3s; in-plane resolution, 3mm; interslice gap, 0.7mm. At the same session, an inversion recovery EPI dataset was also acquired from 43 near-axial 3-mm-thick slices parallel to the AC-PC line: TE, 80ms; TI, 180ms; TR, 16s; in plane resolution 1.5mm; number of signal averages = 8.

Periodic change in T_2^* -weighted signal intensity at the (fundamental) experimentally determined frequency of alternation between A and B conditions (= 1/60 Hz) was

modelled by the sum of a sine wave and cosine wave at that frequency. The amplitudes of the sine and cosine waves, γ and δ respectively, were estimated by pseudogeneralized least-squares fit to the movement-corrected time fMRI series at each voxel. The sum of squared amplitudes, γ^2 and δ^2 , divided by its standard error, provided a standardised estimate of experimentally determined power, the fundamental power quotient (FPQ) (Bullmore et al., 1996). The sign of γ indicated the phase of periodic signal change with respect to the input function. Maps were constructed to represent FPQ and γ at each voxel of each observed dataset. Each observed time series was randomly permuted 10 times, and FPQ estimated as above in each randomized time series, to generate 10 randomized parametric maps of FPQ for each subject in each anatomical plane.

To construct generic brain activation maps, observed and randomized FPQ maps derived from each subject were transformed into the standard space of Talairach and Tournoux and smoothed by a 2D Gaussian filter (SD = 4.5mm) (Talairach & Tournoux, 1988). The median value of FPQ at each intracerebral voxel in standard space was then tested against a critical value of the randomisation distribution for median FPQ ascertained from the randomised FPQ maps. For a one-tailed test of size $\alpha = 0.0008$, the critical value was the $100 \times (1-\alpha)$ th percentile value of the randomisation distribution. Maps of γ observed in each individual were likewise transformed into standard space and smoothed. The median value of γ was computed for each generically activated voxel. If median $\gamma > 0$, that voxel was considered to be generically activated by the gender recognition task (A); if median $\gamma < 0$, that voxel was considered to be generically activated by the theory of mind task (B).

To estimate the difference between control and autism groups in the mean power of response to task B, we fitted the following ANOVA model at each of 1,658 voxels generically activated by the ToM task in one or both of the groups:

$$FPQ_{i,j} = \mu + \beta_1 \text{Group}_j + \varepsilon_{i,j}.$$

Here, $FPQ_{(i,j)}$ denotes the standardized power of response at the i th individual in the j th group. Group denotes a factor coding the main effects of diagnostic status. The null hypothesis of zero between-group difference in mean FPQ was tested by comparing the observed coefficient β_1 to critical values of its non-parametrically ascertained null distribution. To do this, the elements of Group were randomly permuted 10 times at each voxel; β_1 was estimated at each voxel after each permutation; and these estimates were pooled over all intracerebral voxels in standard space to sample the randomisation distribution of β_1 (Brammer et al., 1997). Critical values for a two-tailed test of size $\alpha = 0.01$ were the $100 * (\alpha/2)$ th and $100 * (1-\alpha/2)$ th percentiles of this distribution (Edgington, 1980). For this size of test ($\alpha=0.01$) and search volume (1,658 voxels), we expect no more than 16 voxels to be Type I (false positive) errors⁴.

An analysis of variance in this context assumes that it is meaningful to characterise pathological differences in functional activation in terms of a quantitative difference in mean power of response at each voxel. This

⁴ In fact, 51 voxels were found to have significantly greater power of response to the theory of mind task in controls compared to autistics; and 7 voxels had significantly greater power of response in autistics compared to controls.

assumption has been widely adopted in previous functional imaging studies of neuropsychiatric disorder, most notably it is central to characterisation of schizophrenic abnormalities of functional anatomy in terms of hypofrontality (Weinberger & Berman, 1998). There is also evidence from previous imaging studies of normal subjects that the magnitude of functional response in a given region may be proportional to the cognitive processing load imposed by experimental design (eg (Price & Friston, 1997; Price, Moore & Frackowiak, 1996)). It therefore seems reasonable to interpret differences in power of functional response between control and patient groups as a proxy measure of differences in local neural processing which reflect differences in cognitive strategy imposed by disease.

Results

Considering task performance, both the autism and normal control groups performed both tasks significantly better than chance during scanning. The control group was more accurate in both gender recognition ($x = 86\%$, $sd = 3.0$) and theory of mind ($x = 83\%$, $sd = 7.3$) than the autism group ($x = 82\%$, $sd = 7.5$ and $x = 74\%$, $sd = 1.8$ correct, respectively). For both tasks, there was a significant effect of group, with the normal controls performing better than the subjects with autism or AS (analysis of variance [ANOVA], theory of mind: $F(1,16) = 6.1$, $p = 0.02$; Gender recognition: $F(1,16) = 15.6$, $p = 0.001$).

Functional MRI data were analysed in two stages: First, generic brain activation maps were constructed separately for the control and autism groups. These maps identified

voxels demonstrating significant *power* of periodic signal change over all subjects in each group; they also represented differences between generically activated voxels in terms of *phase* of response to the experimental input function. Thus it was possible to determine which voxels were activated in each group by each of the two tasks.

Second, we used ANOVA to identify voxels that demonstrated a significant difference between groups in mean power of response to each task (see Methods).

Figure 2 shows the functional system activated by presentation of the theory of mind task in the control and autism groups. This system can be anatomically subdivided into two main components: (i) a set of *fronto-temporal* neocortical regions, comprising left dorsolateral prefrontal cortex (DLPFC) approximately Brodmann area (BA) 44, 45, 46; the left medial frontal cortex MFC (BA 9); supplementary motor area (SMA) (medial BA 6); and bilateral temporo-parietal regions, including middle and superior temporal, angular and supramarginal gyri (BA 21, 22, 39, and 40); and (ii) a number of *non-neocortical* areas, including the left amygdala, the left hippocampal gyrus (BA 27 and 30), bilateral insulae, and left striatum.

The autism group activated the frontal components less extensively than the control group; and did not activate the amygdala at all. As shown in Table 2, the control group demonstrated significantly greater power of response in the left amygdala, right insula, and left inferior frontal gyrus. The autism group demonstrated significantly greater power of response in bilateral superior temporal gyrus (STG).

insert Table 2 and Figure 2 here

Discussion

These results are a striking confirmation of Brothers' theory that extracting socially relevant information from visual stimuli is normally associated with activation of the STG, areas of prefrontal cortex⁵, and the amygdala. We next discuss each of these neural regions in turn, both in relation to normal functioning, and to autism.

Regarding the left amygdala, this area may be critically involved in identifying mental state/ emotional information from complex visual stimuli such as the eye region. This laterality effect is consistent with previous studies: the left amygdala appears to be specifically activated in emotion processing (Ketter et al., 1996; Morris et al., 1996)[but see (Breiter et al., 1996; Phillips et al., 1997)]. The autism group appear not to perform the task using the amygdala, but instead place a greater processing load on temporal lobe structures, specialized for verbally labelling complex visual stimuli and processing faces and eyes. We interpret this as showing that people with autism may be solving the task using both language and facial memory functions, perhaps in compensation for an amygdala abnormality. Although it is known that the amygdala plays a role in the recognition of fear (Adolphs et al., 1994; Calder et al., 1996; Scott et al., 1997; Young et al., 1996), here we have also shown that it is involved in inference of a broader range of mental states, from the face and especially the eyes. We consider it unlikely that these results simply reflect emotion-processing or arousal, as the stimuli in the present study involve judging expressions of a broad range of mental states, many of which are not primarily

⁵ We have no strong evidence for OFC activation in these data. This may reflect magnetic susceptibility artefacts induced by the proximity of frontal bone and air spaces.

emotional (e.g., interest, reflective, ignoring). Furthermore, whereas previous studies showing amygdala activation have involved passive perception of powerful emotional stimuli, our task involved an active judgement of a different kind: attribution of a mental state. This suggests that mental state concepts are processed in this region, both when the task involves inferring these from eyes, or other animate actions (Bonda, Petrides, Ostry & Evans, 1996).

Regarding the left prefrontal regions, these may subserve the verbal working memory/central executive function (D'Esposito et al., 1995; Frith, Friston, Liddle & Frackowiak, 1991; Salmon et al., 1996), entailed in matching words whilst observing the eyes. A previous study of autism suggested attenuated activation of MFC (Happé et al., 1996). In the present study we also found MFC was activated less extensively by the autism group, but this was not statistically significant.

Regarding the temporal regions, these may be involved in the **processing** of words and serve as a word store (Wise et al., 1991), and in the processing of eyes (Perrett et al., 1990). We consider that the STG activation seen in the theory of mind condition here is likely to reflect the processing of eyes and faces since it involved bilateral activity, whereas the processing of words would be more likely to have only activated STG lateralized to the left side.

The fundamental premise of this study is that social intelligence is modular or dissociable from general intelligence. More specifically, we have assumed that it will be possible to design a periodic contrast between two experimental conditions which differ exclusively in terms of social cognition, and that the

experimental response to this design will be specific to elements of the "social brain". The validity of this set of assumptions is supported by the correspondence between our results in normal subjects and a prior model of the social brain (Brothers & Ring, 1992). But it may be instructive also to note some limitations and ambiguities inherent in our design.

The two contrasting conditions, although closely matched for stimulus frequency and motor response, may not have differed exclusively in terms of social cognition. For example, subjects may have attempted to solve the experimental problem of mental state assignment by retrieval from long term memory, or by inducing in themselves the emotional states represented by the stimuli. The theory of mind task involved presentation of novel word pairs with each set of visual stimuli, whereas the same pair of short, high frequency words ("male" and "female") were repeatedly presented with each set of stimuli during the gender assignment task. It is thus possible that the experimental contrast could have caused periodic signal change in areas specialised for novelty detection, or differential engagement of languagesystems. Finally, the simultaneous presentation of visual and verbal stimuli, although necessary so that response during scanning could be monitored by forced choice button press, allows an important ambiguity: Do subjects match the eyes to associations or memories primarily induced by the words, or vice versa? In short, the design does not allow us to implicate a particular modality of stimulation (visual or verbal) in experimental activation of the social brain. Several of these problems are typical of periodic or subtraction designs generally, and it will be important in future work to consider so-called parametric experimental designs, in which a single

task is presented at continuously variable levels of difficulty during fMRI data acquisition.

Abnormalities of functional activation by patient groups have often been attributed simply to failure of the patients to perform the task. This seems an inadequate explanation of our findings since the patients performed both tasks better than chance during scanning and had no difficulty in comprehending examples of mental state adjectives presented to them before scanning. However, a number of possible interpretations remain open. It could be that patients with autism have a general deficit in emotional processing, rather than specifically emotional processing to inform mental state assignment. Such a possibility is attractive simply because it is known that the amygdala responds to fearful faces (Breiter et al., 1996; Morris et al., 1996), and that such amygdala activity occurs regardless of whether the subject are aware of the face (Whalen et al, 1998) or aware **that different** facial expressions were **critical to the study** (Morris et al., 1996). **However, since we regard emotional processing as part of social intelligence this interpretation is a refinement rather than a contradiction of our preferred interpretation that autistic patients fail to activate the social brain. Furthermore,** whilst this might be part of the explanation, it cannot be sufficient, since some of the expressions were of non-affective mental states (e.g., “reflective”).

A more problematic alternative is that the patients with autism may in fact activate the social brain, but under both experimental conditions. This pattern of response would not engender periodic signal change and cannot be excluded on the basis of these data. However, even if it were true that the subjects with

autism promiscuously activated the social brain under both conditions, this would still constitute interesting evidence for abnormal modularity or modularisation (Karmiloff-Smith, 1992) of social intelligence in autism. (Here we use the term modularity not in the strong Fodorian (Fodor, 1983) sense, but in a weaker sense

(Baron-Cohen, 1994; Baron-Cohen, in press). **Against this**, however, analysis of the individual subject scans in each condition shows little if any evidence of amygdala activity in the volunteers with autism, which renders the amygdala theory of autism quite plausible.

A further alternative account of the present results might be that people with autism have simply had less experience of the relevant mental states or attitudes being expressed towards them. This also seems unlikely, in relation to states such as ‘sympathy’, ‘reflective’, ‘sad thought’, and ‘friendly’. These are not rare sorts of expressions, and there is no reason to expect that others would not have shown such attitudes towards the subjects in both groups equally. Of course, none of these alternative explanations rules out that the subjects with autism might not *understand* such concepts and expressions less well than controls, but that is precisely the hypothesis that was tested.

Three final alternative accounts might be that eye-movements made by subjects with autism during Task B might have differed significantly in comparison to Task A. We cannot see why the stimuli in Tasks A and B might have provoked different patterns of eye-movement/ visual scanning, since the stimuli were identical in both conditions. However, this remains a small possibility since it may be that when one understands a

visual scene less well, that one scans it less. This should be checked in future studies. A vague and untestable account might be that the autistic group simply expend less 'effort' in attempting to solve such tasks. We do not consider this further as this could never be determined, and in any case would not necessarily be independent of a comprehension deficit.

Future studies are also needed since a task like this could be dismantled into multiple, simpler mental elements. First, patients with autism should be presented with the eyes and no words, and vice versa, to establish which neural activations are due to these two separate factors. Secondly, it will be important to attempt to activate the amygdala in these patients, using a range of cognitive paradigms, to test if the present results reflect a general hypofunctioning of this structure, or whether this is specific to tasks involving inferring mental state. Converging evidence from another social intelligence task will also be important, since the above study employs just one such task. But the present study provides strong evidence of the role of the amygdala in normal social intelligence, and abnormality of the amygdala in autism.

Figure 1: Examples of the stimuli used. During Task B photographs of eyes were presented with a choice of mental state words (examples as shown); during Task A the eyes were presented with a choice of the words “male” and “female”. (Top example: Correct word in Task B = Concerned; Correct word in Task A = Female. Bottom example: Correct word in Task B = Sympathetic; Correct word in Task A = Female).

Figure 2: Generic brain activation maps separately computed from the control and autistic group data are superimposed in standard space. Only those voxels with maximum signal during the theory of mind task are shown. Voxel-wise probability of Type I error $\alpha = 0.008$ for both maps. Voxels activated in the control group only are coloured yellow; voxels activated in the autism group only are coloured red; voxels activated coincidentally in both groups are coloured blue. The right side of each map represents the left side of the brain. The z coordinates (mm) of each slice relative to the intercommissural line in the standard space (Talairach & Tournoux, 1988) is shown above or below each slice.

At -7 mm, the control group activated regions including bilateral insulae and left amygdala; at -2 mm, the main focus of activation in the control group is located in left parahippocampal gyrus; at +10 mm, the control group demonstrates activation of bilateral superior temporal gyrus (STG) and left prefrontal cortex, while the autism group demonstrates less extensive activation of predominantly left sided STG; at +26 and +32 mm, both groups activate left prefrontal cortex.

	Autism	Controls
Age (yrs)		
x	26.3	25.5
sd	2.1	2.8
IQ		
x	108.5	110
sd	10.5	8.5
Handedness (R:L)	6:0	12:0

Table 1: Age, IQ, and handedness of subjects in the experiment.

Cerebral region	BA	Side	N (voxels)	<i>x</i>	<i>y</i>	<i>z</i>	Difference	P
Superior Temporal Gyrus*	22	L	12	-55	-28	15	A > C	0.004
Superior Temporal Gyrus	22	R	8	40	-28	15	A > C	0.002
Inferior Frontal Gyrus+	44/45	L	5	-46	22	9	C > A	0.001
Insula		R	5	40	11	-7	C > A	0.001
Amygdala		L	4	-23	-11	-7	C > A	0.001

Table 2: Main brain regions differentially activated by theory of mind task between control (C) and autism (A) groups. BA = Brodmann Area.

* (or Wernicke's area) + (or Broca's area)

References

Adolphs, R., Tranel, D., Damasio, H., & Damasio, A. (1994). Impaired recognition of emotion in facial expressions following bilateral damage to the human amygdala. Nature, *372*(669-672).

APA. (1994). DSM-IV Diagnostic and Statistical Manual of Mental Disorders, 4th Edition. Washington DC: American Psychiatric Association.

Bachevalier, J. (1991). An animal model for childhood autism: memory loss and socioemotional disturbances following neonatal damage to the limbic system in monkeys. In C. Tamminga & S. Schulz (Eds.), Advances in Neuropsychiatry and Psychopharmacology: Volume 1. Schizophrenia Research, . New York: Raven Press.

Baron-Cohen, S. (1994). How to build a baby that can read minds: Cognitive mechanisms in mindreading. Cahiers de Psychologie Cognitive/ Current Psychology of Cognition, *13*, 513-552.

Baron-Cohen, S. (in press). Does the study of autism justify minimalist innate modularity? Learning and Individual Differences.

Baron-Cohen, S., & Hammer, J. (1997). Parents of children with Asperger Syndrome: what is the cognitive phenotype? Journal of Cognitive Neuroscience, *9*, 548-554.

Baron-Cohen, S., Jolliffe, T., Mortimore, C., & Robertson, M. (1997). Another advanced test of theory of mind: evidence from very high functioning adults with autism or Asperger Syndrome. Journal of Child Psychology and Psychiatry, *38*, 813-822.

Baron-Cohen, S., & Ring, H. (1994). A model of the mindreading system: neuropsychological and neurobiological perspectives. In P. Mitchell & C. Lewis (Eds.), Origins of an understanding of mind, : Lawrence Erlbaum Associates.

Baron-Cohen, S., Ring, H., Moriarty, J., Shmitz, P., Costa, D., & Ell, P. (1994). Recognition of mental state terms: a clinical study of autism, and a functional neuroimaging study of normal adults. British Journal of Psychiatry, 165, 640-649.

Baron-Cohen, S., Wheelwright, S., Stone, V., & Rutherford, M. (submitted). A mathematician, a physicist, and a computer scientist with Asperger Syndrome: performance on folk psychology and folk physics test. .

Bauman, M., & Kemper, T. (1988). Limbic and cerebellar abnormalities: consistent findings in infantile autism. Journal of Neuropathology and Experimental Neurology, 47, 369.

Bauman, M., & Kemper, T. (1994). The Neurobiology of Autism. Baltimore: Johns Hopkins.

Bolton, P., & Griffiths, P. (1997). Association of tuberous sclerosis of temporal lobes with autism and atypical autism. The Lancet, 349, 392-395.

Bonda, E., Petrides, M., Ostry, D., & Evans, A. (1996). Specific involvement of human parietal systems and the amygdala in the perception of biological motion. Journal of Neuroscience, 15(3737-3744).

Brammer, M., Bullmore, E., Simmons, A., Williams, S., Grasby, P., Howard, R., Woodruff, P., & Rabe-Hesketh, S. (1997). Generic brain activation mapping in fMRI: a non-parametric approach. Magnetic Resonance Imaging, 15, 763-770.

Breiter, H. C., Etcoff, N. L., Whalem, P. J., Kennedy, W. A., Rauch, S. L., Buckner, R. L., Strauss, M. M., Hyman, S. E., & Rosen, B. R. (1996). Response and

habituation of the human amygdala during visual processing of facial expression. Neuron, *17*, 875-887.

Brothers, L. (1990). The social brain: a project for integrating primate behaviour and neurophysiology in a new domain. Concepts in Neuroscience, *1*, 27-51.

Brothers, L., & Ring, B. (1992). A neuroethological framework for the representation of minds. Journal of Cognitive Neuroscience, *4*, 107-118.

Brothers, L., Ring, B., & Kling, A. (1990). Responses of neurons in the macaque amygdala to complex social stimuli. Behavioural Brain Research, *41*, 199-213.

Bullmore, E., Brammer, M., Williams, S., Rabe-Hesketh, S., Janot, N., David, A., Mellers, J., Howard, R., & Sham, P. (1996). Statistical methods of estimation and inference for functional MR image analysis. Magnetic Resonance in Medicine, *35*, 261-277.

Calder, A., J, Young, A., W, Rowland, D., Perrett, D., I, Hodges, J., R, & Etcoff, N., L. (1996). Facial emotion recognition after bilateral amygdala damage: Differentially severe impairment of fear. Cognitive Neuropsychology, *13*, 699-745.

Campbell, R., Heywood, C., Cowey, A., Regard, M., & Landis, T. (1990). Sensivity to eye gaze in prosopagnosic patients and monkeys with superior temporal sulcus ablation. Neuropsychologia, *28*, 1123-1142.

D'Esposito, M., Detre, J. A., Alsop, D. C., Shin, R. K., Atlas, S., & Grossman, M. (1995). The neural basis of the central executive system of working memory. Nature, *378*, 279-281.

- Damasio, A., Tranel, D., & Damasio, H. (1990). Individuals with sociopathic behaviour caused by frontal lobe damage fail to respond autonomically to socially charged stimuli. Behavioural Brain Research, *14*, 81-94.
- Edgington, E. S. (1980). Randomisation Tests. New York: Marcel Dekker.
- Eslinger, P., & Damasio, A. (1985). Severe disturbance of higher cognition after bilateral frontal lobe ablation: Patient EVR. Neurology, *35*, 1731-41.
- Fletcher, P. C., Happe, F., Frith, U., Baker, S. C., Dolan, R. J., Frackowiak, R. S. J., & Frith, C. D. (1995). Other minds in the brain: a functional imaging study of "theory of mind" in story comprehension. Cognition, *57*, 109-128.
- Fodor, J. (1983). The modularity of mind: MIT/Bradford Books.
- Frith, C., Friston, K., Liddle, P., & Frackowiak, R. (1991). A PET study of word finding. Neuropsychologia, *29*, 1137-1148.
- Gillberg, I., Bjure, J., Uvebrant, P., Vestergren, E., & Gillberg, C. (1993). SPECT in 31 children and adolescents with autism and autistic like syndromes. European Child and Adolescent Psychiatry, *2*, 50-59.
- Goel, V., Grafman, J., Sadato, N., & Hallett, M. (1995). Modeling other minds. Neuroreport, *6*, 1741-1746.
- Happe, F., Ehlers, S., Fletcher, P., Frith, U., Johansson, M., Gillberg, C., Dolan, R., Frackowiak, R., & Frith, C. (1996). "Theory of mind" in the brain. Evidence from a PET scan study of Asperger Syndrome. NeuroReport, *8*, 197-201.
- Humphrey, N. (1984). The social function of the intellect, N. Humphrey Consciousness Regained, (pp. 14-28): Oxford University Press .
- Karmiloff-Smith, A. (1992). Beyond modularity: MIT Press/Bradford Books.

Karmiloff-Smith, A., Grant, J., Bellugi, U., & Baron-Cohen, S. (1995). Is there a social module? Language, face-processing and theory of mind in William's Syndrome and autism, in press. Journal of Cognitive Neuroscience, *7*, 196-208.

Ketter, T., Andreason, P., George, M., Lee, C., Gill, D., Parekh, P., Willis, M., Herscovitch, P., & Post, R. (1996). Anterior paralimbic mediation of procaine induced emotional and psychosensory experience. Archives of General Psychiatry, *53*, 59-69.

Kling, A., & Brothers, L. (1992). The Amygdala and social behavior. In J. Aggleton (Ed.), Neurobiological aspects of emotion, memory, and mental dysfunction, . New York: Wiley-Liss, Inc.

Morris, J., Frith, C., Perrett, D., Rowland, D., Young, A., Calder, A., & Dolan, R. (1996). A differential neural response in the human amygdala to fearful and happy facial expressions. Nature, *383*(812-815).

Ogawa, S., Lee, T., Kay, A., & Tank, D. (1990). Brain magnetic resonance imaging with contrast dependent blood oxygenation. Proceedings of the National Academy of Science (USA), *3*, 9868-9872.

Perrett, D., Harries, M., Mistlin, A., Hietanen, J., Benson, P., Bevan, R., Thomas, S., Oram, M., Ortega, J., & Brierley, K. (1990). Social signals analyzed at the single cell level: someone is looking at me, something touched me, something moved! International Journal of Comparative Psychology, *4*, 25-55.

Perrett, D., Smith, P., Potter, D., Mistlin, A., Head, A., Milner, A., & Jeeves, M. (1985). Visual cells in the temporal cortex sensitive to face view and gaze direction. Proceedings of the Royal Society of London, *B223*, 293-317.

Phillips, M., Young, A., Senior, C., Brammer, M., Andrew, C., Calder, A., Bullmore, E., Perrett, D., Rowland, D., Williams, S., Gray, J., & David, A. (1997). A

specific neural substrate for perceiving facial expressions of disgust. Nature, 389, 495-498.

Price, C., & Friston, K. (1997). The temporal dynamics of reading: a PET study. Proceedings of the Royal Society, 264, 1785-1791.

Price, C., Moore, C., & Frackowiak, R. (1996). The effect of varying stimulus rate and duration on brain activity during reading. NeuroImage, 3, 40-52.

Rapin, I., & Katzman, R. (1998). Neurobiology of Autism. Annals of Neurology, 43, 7-14.

Sacks, O. (1994). An anthropologist on Mars.

Salmon, E., Van der Linden, M., Collette, F., Maquet, P., Degueldre, C., Luxen, A., & Franck, G. (1996). Regional brain activity during working memory tasks. Brain, 119, 1617-1625.

Scott, S., Young, A., Calder, A., Hellawell, D., Aggleton, J., & Johnson, M. (1997). Impaired auditory recognition of fear and anger following bilateral amygdala lesions. Nature, 385, 254-257.

Tager-Flusberg, H., Boshart, J., & Baron-Cohen, S. (1998). Reading the windows of the soul: evidence of domain specificity sparing in Williams syndrome. Journal of Cognitive Neuroscience, 10, 631-639.

Talairach, J., & Tournoux, P. (1988). Coplanar stereotaxic atlas of the human brain. New York: Thieme Medical Publishers.

Tranel, D., & Hyman, B., T. (1990). Neuropsychological correlates of bilateral amygdala damage. Arch Neurol, 47, 349-355.

Weinberger, D., & Berman, K. (1998). Prefrontal function in schizophrenia: confounds and controversies. In A. Roberts, T. Robbins, & L. Weiskrantz (Eds.),

The Prefrontal Cortex: Executive and cognitive functions, . Oxford: Oxford University Press.

Whiten, A. (1991). Natural theories of mind. Oxford: Basil Blackwell.

Wise, R., Chollet, F., Hadar, U., Friston, K., Hoffner, E., & Frackowiak, R. (1991). Distribution of cortical neural networks involved in word comprehension and retrieval. Brain, 114, 1803-1817.

Young, A., Hallowell, D., De Wal, C., & Johnson, M. (1996). Facial expression processing after amygdectomy. Neuropsychologia, 34, 31-39.

Overleaf: Tables 3 and 4: These are provided for completeness, and show the full results of our preliminary study. Table 3 shows all the regions activated by the ‘theory of mind’ (or social intelligence) condition in the control group, and Table 4 shows all the regions activated by the same condition but in the autism group.

Table 3

Size	Tal(x)	Tal(y)	Tal(z)	Side	Brodmann's Area	Region
169	-43	25	4	L	45	inferior frontal gyrus
163	6	6	53	R	6	medial frontal gyrus
142	43	-33	4	R	21	middle temporal gyrus
122	-17	3	-2	L		putamen
111	-43	3	42	L	6	middle frontal gyrus
101	-46	14	31	L	44	inferior frontal gyrus
82	-23	-6	59	L	6	precentral gyrus
75	-55	-39	9	L	22	superior temporal gyrus
73	32	-72	4	R	19	middle occipital gyrus
66	-49	11	20	L	6	inferior frontal gyrus
63	-3	-44	53	L	7	precuneus
60	-46	14	37	L	9	middle frontal gyrus
50	0	44	15	R	32	cingulate gyrus
42	49	14	26	R	44	inferior frontal gyrus
42	52	-50	15	R	22	superior temporal gyrus
37	38	6	37	R	6	middle frontal gyrus
33	-14	-31	-2	L	27	lingual gyrus
32	0	-47	59	R	7	precuneus
29	-3	-44	37	L	31	cingulate gyrus
26	58	8	15	R	6	inferior frontal gyrus
26	-32	-56	42	L	19	inferior parietal lobule
25	6	3	42	R	24	cingulate gyrus
23	0	-33	31	R	31	cingulate gyrus
22	40	-58	20	R	39	superior temporal gyrus
22	-14	-11	15	L		thalamus
21	-46	-44	26	L	40	supramarginal gyrus
17	29	-58	-7	R	19	fusiform gyrus
16	-46	-22	37	L	1	postcentral gyrus
15	49	-53	-2	R	37	inferior temporal gyrus
15	32	-17	4	R	72	insula
14	23	-56	-13	R		cerebellum
14	12	-72	-2	R	18	lingual gyrus
13	0	47	9	R	32	medial frontal gyrus
13	-29	39	20	L	10	superior frontal gyrus
12	46	11	-7	R	38	superior temporal gyrus
10	52	19	20	R	45	inferior frontal gyrus
10	-3	-86	4	L	18	lingual gyrus
9	-26	-11	-7	L		amygdala
8	-26	-67	31	L	19	angular gyrus
8	-38	-44	-7	L	36	fusiform gyrus
8	35	-22	53	R	4	precentral gyrus
8	29	-22	-2	R		putamen

8	-35	-25	15	L	42	superior temporal gyrus
7	-3	36	-2	L	24	cingulate gyrus
7	20	-53	-2	R	19	lingual gyrus
7	-32	-56	48	L	7	superior parietal lobule
6	52	6	9	R	6	precentral gyrus
6	0	-31	4	R		thalamus
5	-49	-56	15	L	39	superior temporal gyrus
4	-14	-78	-13	L		cerebellum
4	-35	-17	9	L	72	insula
4	-23	-44	59	L	7	postcentral gyrus
1	20	-8	-7	R		amygdala

Table 4

N (Voxels)	x	y	z	Side	BA	Cerebral Region
28	14	-72	-13	R	71	cerebellum
8	17	-53	9	R	23	cingulate gyrus
7	3	-33	31	R	31	cingulate gyrus
15	3	19	31	R	32	cingulate gyrus
30	20	-81	9	R	18	cuneus
9	26	-42	-13	R	36	fusiform gyrus
65	-46	3	26	L	6	inferior frontal gyrus
8	-43	14	20	L	45	inferior frontal gyrus
16	-38	28	20	L	46	inferior frontal gyrus
5	-49	-17	-2	L	21	insula
8	-23	-75	-2	L	18	lingual gyrus
23	-9	50	20	L	9	medial frontal gyrus
37	-43	8	37	L	6	middle frontal gyrus
24	-40	6	42	L	9	middle frontal gyrus
16	-38	28	26	L	46	middle frontal gyrus
6	-23	-75	4	L	18	middle occipital gyrus
16	-49	-42	9	L	21	middle temporal gyrus
8	-43	-58	26	L	39	middle temporal gyrus
8	40	-31	59	R	1	postcentral gyrus
4	-43	-11	42	L	4	precentral gyrus
8	-35	-11	59	L	6	precentral gyrus
9	-14	-44	53	L	7	precuneus
63	6	-53	59	R	7	precuneus
43	-46	-39	15	L	22	superior temporal gyrus
9	43	-31	15	R	22	superior temporal gyrus
19	-49	-17	9	L	42	superior temporal gyrus
6	55	-11	9	R	42	superior temporal gyrus
4	-52	-47	26	L	40	supramarginal gyrus
5	55	-39	26	R	40	supramarginal gyrus